



**Cystatin C** measurements are used as an aid in the diagnosis and treatment of **renal disease**.

METHOD: COLLOIDAL GOLD METHODOLOGY

FEATURES:

FOR THE QUANTITATIVE MEASUREMENT OF CYSTATIN C  
**Genzyme Cystatin C Reagent**

The Genzyme Cystatin C assay uses a colloidal gold methodology that is unique to the cystatin C marketplace and is intended for the measurement of cystatin C concentration in human serum, heparinized plasma, or EDTA plasma.

- ✓ Liquid ready to use reagent and calibrator
- ✓ Analytical sensitivity and excellent precision
- ✓ Extensive list of interference testing completed
- ✓ Applicable to multiple chemistry platforms
- ✓ Serum, EDTA plasma and heparinized plasma acceptable

BENEFITS:

- ✓ Easy to use, no additional preparation required
- ✓ Accurate results
- ✓ High reliability of testing
- ✓ Laboratory flexibility
- ✓ Flexible sample types to meet different laboratory needs

ORDERING INFORMATION	CONFIGURATION	CATALOG NUMBER
<input type="radio"/> GENZYME CYSTATIN C REAGENT	R1 2 X 20 mL R2 2 X 5 mL	80-7136-00
<input type="radio"/> GENZYME CYSTATIN C REAGENT	R1 2 X 48 mL R2 2 X 12 mL	80-7143-00
<input type="radio"/> GENZYME CYSTATIN C CALIBRATOR	6 X 1.5 mL	80-7149-00



**THE AMERICAS**  
 31 New York Avenue  
 Framingham, MA 01701-9322  
 Phone: 800 332 1042  
 Fax: 800 762 6311  
 Email: usdiagcustomerservice@genzyme.com

**INTERNATIONAL**  
 50 Gibson Drive  
 Kings Hill, West Malling  
 Kent, ME19 4AF, UK  
 Phone: +44 (0) 1732 220022  
 Fax: +44 (0) 1732 220024/5  
 Email: ukdiagcustomerservice@genzyme.com

