



**Ammonia** measurements are used in the diagnosis of **coma, Reyes Syndrome** and **liver disease**.

METHOD: ENZYMATIC (GLUTAMATE DEHYDROGENASE); ENDPOINT

FEATURES:

FOR THE QUANTITATIVE MEASUREMENT OF AMMONIA

# Ammonia L3K<sup>®</sup>

The Sekisui Ammonia L3K<sup>®</sup> assay uses a spectrophotometric method utilizing glutamate dehydrogenase. It is a convenient method that is intended for the measurement of ammonia in plasma.

- ✓ No significant interference from ascorbic acid or bilirubin
- ✓ One part, stable liquid
- ✓ Applicable to multiple chemistry platforms

BENEFITS:

- ✓ Reduces the need for repeat testing and sample dilutions
- ✓ Easy to use, no additional preparation required
- ✓ Laboratory flexibility

ORDERING INFORMATION

CONFIGURATION

CATALOG NUMBER

○ AMMONIA L3K<sup>®</sup>

R1 2 X 20 mL  
REAGENT  
1 X 15 mL  
CALIBRATOR

293-10



**THE AMERICAS**  
Sekisui Diagnostics, LLC  
4 Hartwell Place  
Lexington, MA 02421  
Phone: 800 332 1042  
Fax: 800 762 6311  
Email: questions@sekisuidiagnostics.com

**INTERNATIONAL**  
Sekisui Diagnostics (UK) Limited  
Liphook Way, Allington,  
Maidstone, Kent, ME16 0LQ, UK  
Phone: +44 (0) 1622 607800  
Fax: +44 (0) 1622 607801  
Email: info@sekisuidiagnostics.com

